


Active Galactic Nuclei: What we would like to know



Chandra image of Cen A



Volker Beckmann
François Arago Centre, APC
Université Paris Diderot
September 9, 2015






Central Engine



- AGN are bright (10^{41} - 10^{48} erg/sec), point-like ($\ll 100$ pc; Woltjer 1959) and persistent
- No stellar emission, nor supernova, etc.
- Accretion onto a massive black hole
- Masses (from reverberation mapping) in the range of 10^4 (NGC 4395) up to $10^{10} M_{\odot}$ (3C 273)
- Accretion
- Angular momentum \rightarrow accretion disk \rightarrow friction \rightarrow heating \rightarrow thermal emission



Artist's impression of an accretion disk
Credits: Owen/Blondin/NCSU



Central Engine



- alpha-disk: viscosity is some factor α times $c_s h$ (c_s = sound speed, h = scale height of disk)
- Radiative efficient: most of the energy is radiated away


- Eddington limit:

$$L_{\text{Edd}} = \frac{4\pi GMm_{\text{pc}}}{\sigma_{\text{T}}} \simeq 1.3 \times 10^{38} \frac{M}{M_{\odot}} \text{ erg s}^{-1}$$

- apparently some black holes accrete radiatively inefficient
- Sgr A*: $2 \times 10^6 M_{\odot}$ and accretion $10^{-5} M_{\odot} \text{ yr}^{-1}$
→ $10^{-9} L_{\text{edd}}$ (Baganoff et al. 2003)

Fabien Casse:

Accretion around compact sources







Kerr Black Holes



- Black holes can have spin
- Swallowing matter with angular momentum makes a spin likely
- Non-rotating (Schwarzschild) black holes behave differently than maximal rotating (Kerr) black holes.
- Blandford-Znajek (1977) mechanism (aka as Penrose process) allows to extract rotational energy from a black hole (dragging of magnetic field lines)
- Important in the context of raising jets

$$j = \frac{cJ}{GM_{\text{BH}}^2}$$





Tchekhovskoy et al. 2012



Kerr Black Holes




How do we proof that a black hole rotates?




Fuerst & Wu (2004)

Newtonian




Special relativity



Transverse Doppler shift


Beaming

General relativity



Gravitational redshift

Line profile



0.5 1 1.5

ν_{obs}/ν_{em}

Fabian et al. (2008)



Kerr Black Holes?



NGC 1365 NuSTAR/XMM-Newton observations:
 $j = 0.97$ (Risaliti et al. 2013, Nature)

Miller & Turner 2013, ApJ 773, L5:
« The Hard X-Ray Spectrum of NGC 1365:
Scattered Light, Not Black Hole Spin »



5. CONCLUSIONS

AGN spectra may contain significant levels of transmitted and scattered continua from partially covering circumnuclear gas, producing the time-invariant “red wing” below the FeK edge and modifying the spectrum above 10 keV. Such gas is already known in NGC 1365. Geometry-dependent Compton scattering must be taken into account, using three-dimensional radiative transfer calculations, when making spectral models and when accounting for the source total luminosity. The expected

Gardner & Done (2013): high spins are rare

Régis Terrier: reflected spectra / Compton+Rayleigh scattering

Philippe Laurent: AGN with Astro-H



Reynolds (2013): all measured spins $j > 0.5$




Unification




Task: find a model with the smallest number of parameters which explains all AGN.

Peterson (1997)




Seyfert 1 spectrum of NGC 5548

Beckmann & Shrader (2012)



Seyfert 2 spectrum of HE 0201-3029





Beckmann & Shrader (2012);
 graphic by Marie-Luise Menzel



Jets in AGN

- Why do some AGN have jets and some don't?
- Strong radio source: synchrotron emission (jet)
- Don't confuse radio quiet (RQ) and radio silent...
- Franceschini et al. (1998): $L_r \propto M_{\text{BH}}^{2.5}$
- Not found in all studies though
- Laor (2000): all quasars with $M_{\text{BH}} > 10^9 M_{\odot}$ are RL
- All $M_{\text{BH}} < 3 \times 10^8 M_{\odot}$ are RQ
- M_{BH} , L/L_{Edd} , inclination angle
- Other effects: smaller black holes accrete at higher Eddington rate
- Luminosity: core only or include jets?
- E.g. Broderick & Fender (2011): Eddington ratio not strongly correlated to RL
- Environment? Black hole spin?
- B-field (Sikora & Begelman 2013) ?
- jets do not scale simply with mass and accretion rate (Gardner & Done 2013)



$$R^* = \left(\frac{f_{5 \text{ GHz}}}{f_{2500}} \right)$$


Bridle et al. (1994)



Jets in AGN



Jets manage to stay confined for >50 kpc




Protons take more energy for acceleration, but could explain better the confinement over long distances and times ($>10^6$ yrs)




Jets in AGN



- Radio galaxies to study morphology of jets
- Blazars show jet physics
- Spectral energy distribution (SED) and variability give:
- Size and density
- Magnetic field
- Doppler factor
- Particle energy distribution



SED of Mrk 421 (Abdo et al. 2011)





Courtesy: Alan Marscher (Boston Univ.)



AGN environment



- There are no “naked” super-massive black holes
- AGN are at the center of galaxies
- Relation between AGN and its host?





Spiral galaxy NGC 4388 (Subaru Telescope, Suprime Cam, NAOJ)



AGN environment




$M_{\text{BH}} [M_{\odot}]$



Semionov & Vasevicius 2006

Gültekin et al. 2009

Velocity dispersion σ [km / s]




- There is no clear correlation with the type of host galaxy
- Massive SMBH sit rather in elliptical galaxies
(but: ellipticals are on average more massive than spirals!)
- Role of bars on the AGN activity not clear (no correlation: Cardamone et al. 2011, correlation: Alonso et al. 2012)



AGN environment



- Feedback between SMBH and host?
- Does the AGN regulate its own feeding?
- Does the AGN regulate the star burst activity in the host?
- Yes: e.g. De Young (2010), Fabian (2010)
- No: Jahnke & Maccio (2011), Jian et al. (2011), Juneau et al. (2012; but intrinsic absorption correlated with star formation rate)
- Gas is needed to feed AGN. 10% of galaxies at $z \sim 2$ are AGN, but only 0.1% at $z=0$ (Dunlop et al. 2003)
- No correlation between AGN axis and host galaxy orientation (Keel 1980, Lawrence & Elvis 1982, Hopkins et al. 2012)





NGC 4151 (Mundell et al. 1999, 2002; Fernandez et al. 1999)



AGN & merger



- Merging of host galaxies transports matter to the core
- Hopkins et al. (2010): uncertainty leaves room for any kind of scenario
- Kocevski et al. (2012): Chandra detected AGN + control sample: major merger do not trigger AGN at $z \sim 2$ (but minor merger?)
- Difference between dry and wet (= gas-rich) merger?
- eLISA will tell us what the real AGN merger rate is (but we might have to wait until 2034...)





SMBH formation



- Quasars observed back to $z=7.085$ (Mortlock et al. 2011)
- How to form a $M > 10^8 M_{\odot}$ black hole within $< 6 \times 10^8$ years ?
- Assume Eddington limited accretion $\lambda=1$, efficiency $\eta = 10\%$:
- $M_0 = 10 M_{\odot} \rightarrow 9 \times 10^6 M_{\odot}$ after 6×10^8 years
- Either lower efficiency, or larger start mass:
- $\eta = 0.09$, $M_0 = 150 M_{\odot} \rightarrow M(z=7) = 6 \times 10^8 M_{\odot}$
- But: accretion at the Eddington limit for 1 billion years?
- Initial mass function unknown
- Quasi stars (low J and metallicity) with direct collapse to $1000 < M < 10^6 M_{\odot}$ (Begelman et al. 2008) ?
- But: galaxies do not look very different at $6 < z < 8.7$ (Mc Lure et al. 2011); similar colours and SFR as at $z \sim 2$
- James Webb Space Telescope to study first stars







SMBH formation



- Other solutions to get a large starting mass:
- Primordial black holes (e.g. Hawkins 2011, but see Pani & Loeb 2014)
- Merging of smaller black holes
- Problems with recoils (e.g. Campanelli et al. 2007) ?
- Starting with a black hole of $M_0 \sim 1000 M_\odot$ it is “easy” to build SMBH until $z \sim 7$ (e.g. simulations by Sijacki et al. 2009)
- eLISA will reveal the merging history in the Universe






Open questions I



- Kerr black holes? Model independent measurement necessary, see **presentation by Philippe**
- Accretion models: see **presentation by Fabien**
- Geometry of AGN? Patchy corona? Better models + observations (X-ray/hard X-ray) necessary
- How to raise the jet? VLBI shows details and starting point
- Related to unknowns in accretion and BH spin
- GBH most luminous when there is no jet -- but AGN are most luminous as radio loud quasars
- GBH: baryons in jet (SS 433 Margon+ 1979 , 4U 1640-47 Diaz Trigo+ 2013)
- What collimates the jet?
- Polarization mapping, higher resolution radio observations, numerical modeling
- **Dimitri: IceCube events from AGN?**








Open questions II



- SED modeling
 - Synchrotron and EC / SSC / mix ?
 - Better sampling needed (and not always the same blazars!)
 - Flaring objects might not be representative
 - Are all non-blazar gamma-ray bright AGN somehow intrinsic blazars? **Santiago: AGN in gamma-rays**
-
- Unification: statistics
 - SDSS, Gaia, eROSITA, LSST, Euclid ... : 10^7 AGN
 - New data treatment techniques needed








Open questions III



- Environment
- Is there feedback or not? Does the host galaxy play a major role (other than providing mass) ? Do bars play a role?
- Merging: important for the AGN? Minor or Major merger?
- better (control) samples needed
- Dark matter in the Universe important
- Euclid, DES, PanSTARRS, PAU, LSST ...
- **Paolo: AGN with XSHOOTER**
- SMBH: how to build a quasar until $z \sim 7$?
- We do not know much about the Universe beyond $z \sim 7$ until CMB
- JWST, E-ELT, theory, numerical simulations
- Absorption evolution? NuSTAR, Astro-H, ATHENA+, eROSITA
- Duty cycle of AGN?





What's needed



- Theoretical models
- Simulations
- Observations
- Processing and data mining

- Get more students and colleagues on this exciting topic!

Beckmann & Shrader 2012,
“Active Galactic Nuclei”, Wiley-VCH,
382 pages
(see e.g. the Condorcet library)

